

Quantum phase transitions in the $J - J'$ Heisenberg and XY spin- $\frac{1}{2}$ antiferromagnets on square lattice: Finite-size scaling analysis

Piotr Tomczak

Physics Department, Adam Mickiewicz University,

Umultowska 85, 61-614 Poznań, Poland

Johannes Richter

Institut für Theoretische Physik, Otto-von-Guericke Universität, Magdeburg,

P.O.B. 4120, 39016 Magdeburg, Germany

Abstract

We investigate the critical parameters of an order-disorder quantum phase transitions in the spin- $\frac{1}{2}$ $J - J'$ Heisenberg and XY antiferromagnets on square lattice. Basing on the excitation gaps calculated by exact diagonalization technique for systems up to 32 spins and finite-size scaling analysis we estimate the critical couplings and exponents of the correlation length for both models. Our analysis confirms the universal critical behavior of these quantum phase transitions: They belong to 3D O(3) and 3D O(2) universality classes, respectively.

The equivalence of the critical behavior of D dimensional quantum spin systems and $D + 1$ dimensional classical spin systems is well recognized. This idea, combined with finite-size scaling was used previously many times to discuss the critical properties of infinite spin systems, see e.g., [1,2] and [3] for a review. However, these investigations were strongly limited with respect to the system size. Due to the recent advances in computer technology it is possible to treat bigger systems, e.g., up to 36 sites for spin $\frac{1}{2}$, and consequently to extract their critical properties using finite-size scaling method [4,5]. Our aim is to present in this paper the results of such an investigation of critical parameters (coupling and exponents of correlation length) for the $J - J'$ Heisenberg and XY spin- $\frac{1}{2}$ antiferromagnets on square lattice.

The Hamiltonian of the model whose critical behavior is examined is given by

$$H = J \sum_{\langle i,j \rangle} \vec{S}_i \cdot \vec{S}_j + J' \sum_{\langle k,l \rangle} \vec{S}_k \cdot \vec{S}_l. \quad (1)$$

The first sum, denoted by $\langle i, j \rangle$, runs over pairs of nearest-neighbors on the square lattice connected by thin bonds (see Fig.1), whereas the second one, denoted by $\langle k, l \rangle$ — over nearest-neighbors connected by thick bonds. In the case of Heisenberg spin system three Pauli matrices are included into scalar product in Eq. (1), in the case of XY system — only two. The model represents an antiferromagnet, i.e., both couplings are positive and additionally, $J' \geq J$. Clearly, what one can see here is the competition between long-range Néel order and the tendency to the formation of local singlets of two neighboring spins, coupled via J' . In limiting cases this model reduces to the long-range ordered Heisenberg antiferromagnet on square lattice for $J = J'$ on one hand and on the other — to disjoint singlets (no staggered magnetization) for $J'/J \rightarrow \infty$. At some (J'/J) there exists a quantum phase transition between the gapless Néel phase and a gapped 'singlet' phase (quantum paramagnet). The properties of $J - J'$ Heisenberg model on the square lattice were first examined by series expansion (SE) [6] and more recently by renormalized spin wave (RSW) approach [7], exact diagonalization (ED) and coupled cluster method (CCM) [9] in order to observe the interplay between the local singlet formation tendency and the long-range

Néel order. Although, in general, all those methods predict the existence of quantum phase transition, they differ drastically in the estimation of the critical coupling. Furthermore, there was only one attempt [6] to find critical exponents, however the error of this estimation was rather large.

Let us now rewrite the Hamiltonian in such a form that the two above tendencies will be seen more clearly:

$$H = g(\sum_{\langle i,j \rangle} \vec{S}_i \cdot \vec{S}_j + \frac{1}{g^2} \sum_{\langle k,l \rangle} \vec{S}_k \cdot \vec{S}_l), \quad (2)$$

$0 < g \leq 1$. Note that in Eq. (2) one has the same summation specification as in Eq. (1). The coupling constant g determines the relevant energy scale in the model under consideration and the term $1/g^2 = \lambda$ is analogous to an inverse temperature.

In what follows the estimation of the critical value of g_c and critical exponent $1/\nu$ for the Heisenberg and XY Hamiltonians will be described. At the beginning one finds by the exact diagonalization (Lanczos algorithm) the spin-gap, defined as $\Delta = E_1 - E_0$ ($E_{0(1)}$ is the lowest energy in the $S^z = 0(1)$ sector) dependence vs. $\lambda = 1/g^2$ for the Hamiltonian (2) on a square lattice for a sequence of finite systems. Note that all the systems, being elements of the sequence should be invariant under the same symmetry operations; in the opposite case it is not possible to find a proper scaling. In the system under examination one has only three such systems: with $N = 8, 18, 32$ spins. Two of them are shown in Fig. 1, the third one has the same shape.

The spin-gap, multiplied by the linear dimension of the system, $\Delta\sqrt{N}$, allows one to find the pseudo-critical points [3]: for the Heisenberg and XY systems one has three such points, collected in Table 1 (Heisenberg) and 2 (XY). Next one should extrapolate the sequence of pseudo-critical points to infinity. However, it is not possible to employ here any algorithm improving the convergence of the finite-lattice data sequence the series is extremely short. Therefore we find it more accurate to use graphical methods to find the critical value of g . The sequences of pseudo-critical points for Heisenberg and XY system plotted vs. $1/L^4$ are shown in Fig. 2. The estimate of $1/g_c^2 = 2.46(2)$ for the infinite Heisenberg system

should be compared to the value 2.56 obtained by SE [6], to the value of 3.16 obtained by CCM method, to the value of 2.45 from ED [9] and finally to the value of 5.0 [7] from RSW approach. The extrapolated value of $1/g_c^2 = 4.56(2)$ for *anisotropic* XY system is higher, as one should expect, than that for Heisenberg system since anisotropy act against the singlet formation [8].

The critical exponent for the correlation length, ν , may be estimated from the behavior of the Callan-Symanzik β -function [10,3,1]

$$\beta(\lambda)/g = \frac{d}{dg} \ln[g\Delta(\lambda)], \quad (3)$$

which, calculated for a finite system of linear size L in a pseudo-critical point, scales as

$$\beta(\lambda_c, L) \sim L^{-1/\nu}. \quad (4)$$

Usually, in order to see this scaling behavior, one takes into account a sequence of $\beta(L_i)/\beta(L_{i-1})$ values calculated at the pseudo-critical points for some values of L_i , such that $L_i = L_{i-1} + 1$. Expanding

$$\beta(L_i)/\beta(L_{i-1}) \sim (1 + 1/L_i)^{-1/\nu} \sim 1 - \frac{1}{\nu} \frac{1}{L_i} + \dots \quad (5)$$

one finds a linear behavior of $(1 - \beta(L_i)/\beta(L_{i-1}))$ vs. $1/L$ for all i . This is the 'linear' approximation. Note that the error is $O(1/L^2)$. However, if one can not find a sequence of finite systems fulfilling $L_i = L_{i-1} + 1$ (in our case $L_i = L_{i-1} + \sqrt{2}$) this approximation is rather crude because of the order of the error. This may be improved in the following way. First, let us note that in the following expansion, for small x

$$\ln\left(\frac{1+x}{1-x}\right) = 2x + \frac{2}{3}x^3 + \dots \quad (6)$$

the term x^2 is absent and the error is $O(x^3)$. Second, let us put

$$x = \frac{L_i - L_{i-1}}{L_i + L_{i-1}}, \quad (7)$$

and expand

$$\ln[\beta(L_{i-1})/\beta(L_i)] \sim \frac{1}{\nu} \ln\left(\frac{1+x}{1-x}\right) \sim \frac{2}{\nu} \frac{(L_i - L_{i-1})}{(L_i + L_{i-1})} + \dots \quad (8)$$

Consequently one has a linear dependence of $\ln[\beta(L_i)/\beta(L_{i-1})]$ vs $1/L$:

$$\ln[\beta(L_{i-1})/\beta(L_i)] \sim \frac{\sqrt{2}}{\nu} \frac{1}{\bar{L}_i} + \dots, \quad (9)$$

where $\bar{L}_i = (L_i + L_{i-1})/2$. The main advantage of this approach is a small finite size error, what in consequence enables one to examine smaller systems. However, there remains yet another problem. It is possible to consider scaling from two pseudo-critical points: (8-18) where $x = 1/5$ and (18-32) — where $x = 1/7$. In the third point the expansion 8 gives rather large finite size correction ($x = 1/3$) and this point has to be excluded. Thus, one can ask whether it is possible to find a scaling relation and to estimate critical exponent from two points only? The answer is yes, but the final error will be larger. Since the expansion (8) produces an error $O(x^3)$, it seems to be especially well suited to this purpose. To test this approach we have extrapolated the $1/\nu$ exponent from the data for transverse Ising model on square lattice [5] taking into account only two pseudo-critical points: (16-25) and (25-36). The estimate of $1/\nu = 1.586(7)$ obtained by simple linear extrapolation from these two points should be compared to the original one $1/\nu = 1.591(1)$ [5], extrapolated from four points.

The data collected in Table 1 and 2 enable one to calculate β -function from Eq. (3) and consequently $1/\nu$ from Eq. (9) for each pseudo-critical point of $J - J'$ model; their values for Heisenberg and XY Hamiltonians are listed in Table 3 and the graphical extrapolation is shown in Fig 3. The values of critical exponents for the same universality classes obtained by other authors are also displayed in Table 3.

One should note that the error in the present approach is larger than in other finite-size scaling analysis, but as was mentioned this is a result of the small number of systems with required symmetry, see Fig. 1, and the error is still comparable with an error resulting from extensive Monte-Carlo simulation [11,12].

To conclude, we have presented the results of the investigation of the critical parameters for the quantum phase transitions in the spin- $\frac{1}{2}$ $J - J'$ Heisenberg and XY antiferromagnets

on square lattice. The obtained values of the correlation length critical exponents strongly suggest that these transitions belong to the 3D $O(3)$ and 3D $O(2)$ universality classes, respectively.

Acknowledgments The authors thank Jörg Schulenburg for the numerical assistance and Prof. Ryszard Ferchmin for reading manuscript. Support from the Polish Committee for Scientific Research (Project No. 2 PO3B 046 14) and from the Deutsche Forschungsgemeinschaft (Projects No. 436/POL17/5/01 and Ri 615/10-1) are also acknowledged. Some of the calculations were performed at the Poznań Supercomputer and Networking Center.

REFERENCES

- [1] C. J. Hamer and Michael N. Barber, J. Phys. A: Math. Gen. **13**, L169 (1980).
- [2] C. J. Hamer, J. Phys. A: Math. Gen. **15**, L675 (1982).
- [3] M. N. Barber, *Phase Transitions and Critical Phenomena* vol. 8, eds. C. Domb and J. L. Lebowitz, Academic, New York (1981).
- [4] C. J. Hamer, Tobias Hövelborn and Michael Bachhuber, J. Phys. A: Math. Gen. **32**, 51 (1999).
- [5] C. J. Hamer, J. Phys. A: Math. Gen. **33**, 6683 (2000).
- [6] Rajiv R. P. Singh, Martin P. Gelfand and David A. Huse, Phys. Rev. Lett **61**, 2484 (1988).
- [7] N. B. Ivanov, S. E. Krüger and J. Richter, Phys. Rev. B **53**, 2633 (1996).
- [8] O. Derzhko, J. Richter, O. Zaburannyi, J. Phys. Condens. Matter **12**, 8661 (2000).
- [9] Sven E. Krüger, Johannes Richter, Jörg Schulenburg, Damian J. J. Farnell, Raymond F. Bishop, Phys. Rev. B **61**, 14607 (2000).
- [10] C. J. Hamer, J. Kogut, Phys. Rev. B **20**, 3859 (1979).
- [11] M. Troyer, M. Imada, and K. Ueda, J. Phys. Soc. Jpn. **66**, 2957 (1997).
- [12] K. Chen, A. M. Ferrenberg and D. P. Landau, Phys. Rev. B **48**, 3249 (1993).
- [13] J. C. Le Guillou and J. Zinn-Justin, Phys. Rev. B **21**, 3976 (1980).
- [14] J. C. Le Guillou and J. Zinn-Justin, J. Physique Lett. **46**, L-137 (1985).
- [15] A. P. Gottlob and M. Hasenbusch, J. Stat. Phys. **77**, 919 (1994).

FIGURES

FIG. 1. The $J - J'$ model on square lattice. Finite systems of 8 and 18 spins are marked by dashed lines.

FIG. 2. Extrapolation of the finite-system pseudo-critical points as a function of $1/L^4$. Open squares - Heisenberg model, filled squares - XY model.

FIG. 3. Extrapolation of the finite-system estimates of the $1/\nu$ exponent as a function of $1/L$. Open squares - Heisenberg model, filled squares - XY model.

TABLES

TABLE I. Pseudo-critical points λ_c calculated for two Heisenberg systems of sizes given in the first column. The values of the gap Δ and its derivative Δ' at the pseudo-critical point λ_c are also listed. To find them 11 ED data points equally spaced around λ_c were fitted to the polynomial of 4-th order in λ in the region of $\lambda_c \pm 0.005$. 5 digits are exact.

System size	λ_c	Δ	Δ'
8	3.1166	4.14495	1.52350
18		2.76330	1.71055
8	2.9251	3.85980	1.45296
32		1.92990	1.81304
18	2.7648	2.20467	1.44663
32		1.65350	1.62565
extrapolated	2.46(2)		

TABLE II. Same as TABLE I, but for XY spin systems.

System size	λ_c	Δ	Δ'
8	4.9326	2.51653	0.786928
18		1.67768	0.848422
8	4.7705	2.39028	0.770058
32		1.19514	0.942901
18	4.6444	1.44152	0.788049
32		1.08114	0.866564
extrapolated	4.56(2)		

TABLE III. Values of critical exponent $1/\nu$ calculated for two pseudo-critical points for finite Heisenberg and XY spin systems and subsequently extrapolated to infinity. For comparison are also included values of the same exponents calculated by other authors.

System size	Heisenberg O(3)	XY, O(2)
8-18	1.9841	1.6220
18-32	1.8323	1.6014
extrapolated	1.44(4)	1.55(4)
	1.46(8) [11]	1.495(5) [13]
	1.418(6) [12]	1.49(1) [14]
		1.51(1) [15]





